



Automated Photometry of γ Cassiopeiae: The Last Roundup

M. A. Smith¹ and G. W. Henry² ¹ NSF's National Optical–Infrared Astronomy Research Laboratory, 950 N. Cherry Ave., Tucson, AZ 85721, USA; myronmeister@gmail.com² Center of Excellence in Information Systems, Tennessee State University, Nashville, TN 37209, USA

Received 2021 January 2; revised 2021 April 29; accepted 2021 May 4; published 2021 June 29

Abstract

γ Cassiopeiae (B0.5IVe) is the noted prototype of a subgroup of classical Be stars exhibiting hard thermal X-ray emission. This paper reports results from a 23 yr optical campaign on this star with an automated photometric telescope (APT). A series of unstable long cycles of length 56–91 days has nearly ceased over the last decade. Also, we revise the frequency of the dominant coherent signal at $0.82238 \text{ days}^{-1}$. This signal's amplitude has nearly disappeared in the last 15 yr but has somewhat recovered its former strength. We confirm the presence of secondary nonradial pulsation signals found by other authors at frequencies 1.24, 2.48, and 5.03 days^{-1} . The APT data from intensively monitored nights reveal rapidly variable amplitudes among these frequencies. We show that peculiarities in the 0.82 days^{-1} waveform exist that can vary even over several days. Although the 0.82 days^{-1} frequency is near the star's presumed rotational frequency, because of its phase slippage with respect to a dip pattern in its far-UV light curve, it is preferable to consider the UV pattern, not the 0.82 days^{-1} signal, as associated with rotational modulation. We also find hints of the UV dip pattern in periodograms of seasonal data early in our program.

Unified Astronomy Thesaurus concepts: B stars (128)*Supporting material:* machine-readable table

1. Introduction

As the prototype of classical Be stars, γ Cassiopeiae (γ Cas; B0.5IVe) has exhibited a long history of multiwavelength variability, ranging from the disappearance of its Be decretion disk to optical and X-ray flaring (Doazan et al. 1983; Harmanec 2002; Smith 2019, “S19”). Apart from its optical variability, it has become an object of interest among high-energy astronomers since its discovery as a hard X-ray emitter (Mason et al. 1976; White et al. 1982). Given these discoveries, γ Cas has been a target of several multiwavelength campaigns.

With a considerable X-ray flux ($L_x/L_{\text{bol}} = 3\text{--}5 \times 10^{-6}$), it is also the prototype of an X-ray class of at least 25 members (Nazé et al. 2020a). These stars are defined by their hard (but thermal) X-ray spectra, which exhibit emission lines from multiple thermal components. The spectrum of γ Cas itself indicates a dominant $kT \approx 14 \text{ keV}$ plasma that overwhelms fluxes at all X-ray energies. Its X-ray light curve is variable over timescales of seconds to more than a year. A peculiar, if not unique, characteristic of the apparently bright members of the class is the existence of ubiquitous rapid X-ray “quasi-flares.” These features have decay times as short as 4 s, proving they are formed in photospheric densities (Smith et al. 1998a, “SRC”). A review of the properties of these unique X-ray Be stars is given in Smith et al. (2016, “SLM16”).

To bridge the optical and X-ray domains, SRC and Smith et al. (1998b, “SRH”) conducted a 34 hr time series analysis with the Short-wavelength Prime (SWP) camera of the International Ultraviolet Explorer (IUE) in 1996 January and simultaneous 21.5 hr monitoring with the Hubble Space Telescope/Goddard High Resolution Spectrometer (GHRS) and the Rossi X-Ray Timing Explorer (RXTE) on 1996 March 14–15. With the decommissionings of the IUE and GHRS, no further UV monitoring has been possible.

Several interesting results came of these campaigns, notably flux correlations between the optical and X-ray domains.

GHRS spectra were binned in wavelength to construct a high signal-to-noise ratio (S/N) and high time resolution quasi-continuum “UVC” light curve. Two prominent ($\sim 1\%$) dips, each lasting a few hours and separated by about 9 hr, were visible. Photospheric UV spectral lines of Si^{2+} , Si^{3+} , Ni^{+} , Fe^{+} , Fe^{4+} , and S^{3+} in the same data set showed correlations/anticorrelations with the UVC curve (Smith & Robinson 1999, “SR99”; Smith & Robinson 2003). These UV diagnostics were in turn correlated with the simultaneous X-ray fluxes. To add to this mix, we found the two dips separated by 9 hr to have been observed by the IUE 57 days earlier. The same dip pattern emerged from IUE observations in 1982 found in the IUE archive. Although the $\lambda\lambda 1407\text{--}1417$ GHRS light curve could not exhibit color changes during the events, the broad wavelength range of the IUE/SWP spectra revealed that color changes were consistent with absorptions by large cool clouds attached to the star over intermediate latitudes (SRH).

Another pattern in the GHRS data was short-lived migrating subfeatures (msf) that moved from blue to red across line profiles. Similar features had been found in optical spectra by Yang et al. (1988) and Smith (1995). Because this phenomenon had been observed only spectroscopically for γ Cas, we were curious to see whether it also has a signature in broadband photometry.

These patterns indicated the need for long-term monitoring of the star. Therefore, in 1997 we initiated a campaign with Tennessee State University's T3 Automated Photometric Telescope (APT). Although the APT shared observing time with several other scientific projects, we planned to observe γ Cas a few times on every available photometric night it was in view. Occasionally we could dedicate full nights to the program.

The first discovery from the APT campaign was of noncoherent, unstable “long cycles” ranging in length from about 56 days to 91 days (Robinson et al. 2002, “RSH”; Smith et al. 2006 (Paper I); Henry & Smith 2012 (Paper II)). The

amplitudes of the cycles observed in V are often larger than those in B , which implies the cycles are caused by density modulations within the decretion disk.

To see if these long cycles were related to the star’s X-ray flux, RSH requested and were granted six 27 hr RXTE visits during 2000–2001. The exposure durations were chosen to average the flux over the star’s estimated rotational period. Intervals between successive visits were doubled, such that RXTE covered a timescale range of a week to almost 11 months. Paper I reported that X-ray and optical fluxes during these intervals revealed a sinusoidal fit to the APT long-cycle variations that, when suitably scaled to the X-ray fluxes, showed a very good match. The authors suggested that the long cycles in the optical and X-ray regimes were caused by a magnetorotational disk dynamo. The authors extended this correlation by demonstrating a reasonably good prediction of the X-ray flux from the ongoing APT monitoring. An updated APT data set was investigated by Motch et al. (2015) using the RXTE All Sky Monitoring data as well as a later Japanese MAXI (X-ray) data set. These authors confirmed the optical/X-ray correlation of Paper I and found that the APT and X-ray data sets correlated from short (a few hours) to very long (weeks or longer) timescales as well. The latter X-ray data sets were independent of the X-ray data in Paper I. Also, these authors found that there is no visible time lag between the two sets of variations, in contrast to lags typical of X-ray Be–neutron star (NS) binaries.

In addition to long cycles, Papers I and II reported a coherent signal with $P = 1.2158$ days ($0.82 \text{ days}^{-1} \equiv f_{82}''$). We noticed that this frequency is consistent with the expected rotational period of ~ 1.24 days, according to our estimated physical parameters for the star. Thus, we adopted 1.21 days as the rotational period. Similarly, we suggested that the periodic UVC dips were caused by the rotational modulation of anchored clouds. This paper will revisit these findings.

2. A Sketch of Suggested X-Ray Mechanisms

As it is relevant to the APT study, we sketch a history of attempts to explain the production of hard, thermal X-ray flux in γ Cas and its association with certain optical and UV variabilities.

Harmanec et al. (2000) discovered that γ Cas is a binary system. Its orbital period is 203.5 days, and it is in a nearly circular orbit ($e \lesssim 0.03$). Although the secondary’s mass is $0.9 \pm 0.1 M_{\odot}$ (Nemravová et al. 2012; Smith et al. 2012, “SLM”), its evolutionary status is unknown. However, the evidence is strong that the Be star is a blue straggler (Mamajek 2017; E. Mamajek 2017, private communication) and therefore probably has an envelope-stripped or degenerate secondary. Assuming the secondary’s mass estimate is accurate, the mass is too low for it to be a NS, but it is appropriate for a white dwarf (WD) and (envelope-stripped) helium stars. The limit straddles the masses of sdO stars.

Wang et al. (2017) have cross-correlated IUE spectra of γ Cas but found no evidence of a far-UV contribution in the far-UV down to a level of 0.6%. This result rules out a range of types of evolved, low-luminosity companions. However, Wang et al. (2021) performed the same tests on 13 other early-type Be stars not previously known to be in binaries and found 10 from this sample exhibit at least traces of a spectrum of a hot subluminal secondary such as an sdO star. Using the same technique, Gies et al. (1998) and Peters et al.

(2008, 2013, 2013) had previously discovered substantial UV contributions from the sdO secondaries of three Be binaries (ϕ Per, FY CMa, and 59 Cyg). In a fourth case, HR 2142, only a faint contribution can be seen from an “sdO in transition.” Clearly, sdO stars are the most likely kind of secondary in early Be systems. Thus, to see if these sdO stars could be seen against γ Cas as the primary star, we substituted the appropriate physical parameters of γ Cas for the parameters of the actual primaries and recomputed the secondary flux contributions to simulated γ Cas–sdO systems. This exercise confirmed that sdO contributions would still be recognizable for the first three imaginary cases if γ Cas had been the primary. We add finally that at least two of the binaries in the Peters–Gies et al. sample, 59 Cyg and ϕ Per (those with the brightest and most massive hot secondaries), are faint and soft X-ray systems (HEASARC Rosat_Data_Archive 2020; Nazé et al. 2020a), which clearly cannot be confused with γ Cas star emissions.

Even before the binarity of γ Cas was discovered, accretion of Be wind onto a degenerate object was suggested as the source of the hard X-ray emission. White et al. (1982) and Murakami et al. (1986) had argued that a secondary ought to be an NS or WD. More recently, various authors have again suggested accretion involving degenerate or hot secondaries: a WD (Hamaguchi et al. 2016), an NS in the propeller stage (Postnov et al. 2017), or interactions between an sdO wind and the Be disk (Langer et al. 2020, “L20”).

The difficulties described by SLM16 in reconciling the unique X-ray characteristics of γ Cas with those of other X-ray classes of Be stars motivated SRC, SR99, and Robinson & Smith (2000, “RS00”) to advance a very different mechanism for the hard X-ray production: the star–disk magnetic interaction hypothesis.

This idea requires the tangling of field lines from putative small-scale magnetic surface complexes (e.g., Cantiello & Braithwaite 2011) and a toroidal field in the inner decretion disk. The different angular rotation rates of the star and the disk cause interactions of star–disk fields, which in a short time entangle, break, reconnect, and ultimately relax. This process releases magnetic energy, which accelerates embedded particles in high-energy beams. Some of these are guided by protruding surface field lines toward the star. In fact, the existence of downstreaming matter can be inferred from highly redshifted absorption lines in the 1996 March GHRS time series (SR99). According to simulations by RS00, nearly monoenergetic (200 keV) electron beams impact and thermalize the surface, causing local explosions (also called “flares”). Their detritus accumulates in low-density canopies and decays in ~ 20 minutes, producing a basal flux of the same high temperature.

3. Observations

We have been conducting Johnson B and V photometric observations of γ Cas since 1997 with the T3 0.4 m APT facility at Fairborn Observatory in southern Arizona (Henry 1995a, 1995b, 1999; Eaton et al. 2003). The APT acquires successive brightness measurements of individual target stars with a single-channel photometer using a temperature-stabilized EMI 9924B photomultiplier tube. Each observation of γ Cas, which we refer to as a group observation, consists of the mean of three B and V differential measurements of γ Cas with respect to its comparison star (HD 6210) and the

mean of two measurements of the check star (HD 5395) with respect to the comparison star.

To avoid saturating the detector on 2nd mag γ Cas, we used a 3.8 mag neutral density filter to observe all three stars, with the exception of the 1997 season, when we used different neutral density filters for γ Cas and the comparison stars.

γ Cas comes to opposition in early October, and our APT observing seasons always covered the months of September through early February. Telescope closure was generally forced by Arizona's summer monsoons. However, if the monsoons arrived late, a new season included some nights in June–July. In the analysis that follows, we label each season by the year corresponding to the star's opposition. Our final data set covers 23 consecutive observing seasons from 1997 through 2019.

On most clear nights the APT was programmed to acquire one to four observations spaced by ≥ 2 hr. During most observing seasons, a few nights near opposition were dedicated to monitoring for several hours. On those nights the APT acquired one to four group observations spaced about 8 minutes apart. On good nights the external precision of the group means was typically 0.003–0.004 mag, as determined from observations of pairs of constant stars. Since the APT can acquire observations in marginal photometric conditions, we rejected as outliers any group mean differential magnitudes with standard deviations greater than 0.01 mag. Because we used different neutral density filters for the variable and comparison stars in the 1997 season, we were forced to adjust those means to the 1998 season means. Finally, the check minus comparison star differential magnitudes demonstrated that both are constant to ≤ 0.005 mag on seasonal and year-to-year timescales.

In this paper we present our final 1997–2019 data set for γ Cas, consisting of 5554 observations in the B passband and 5488 in the V passband. We have cleaned the data set by fitting sine curves to the long-cycle (~ 70 –80 day) variability in single observing seasons and rejected as outliers those observations with residuals from the least-squares sine fit of $\geq 2.5\sigma$. For several of the later observing seasons, where we could not obtain a reliable long cycle, we rejected observations that were $\geq 3.0\sigma$ from the seasonal mean magnitudes.

Our analysis technique employed the frequency search method of Vaníček (1971), based on least-squares fitting of sine curves, to search for periodicities in various combinations of the yearly photometric data sets. This method uses the reduction factor in the data's variance as a goodness-of-fit parameter. For all analyses we initially scanned trial frequencies over a range of 0.005–6.0 days $^{-1}$. Formal uncertainties in the best-fit periods and amplitudes were computed from standard propagation formulae. We consider a signal to be real only if it is found in both B and V data sets. For multiseasonal analyses we forced the seasonal means to the same value to eliminate those low-frequency variations that can be seen in the long-term light curve (Figure 1). If $>3\sigma$ outliers were present at any of the frequencies detected within the individual observing seasons, we removed those observations from our data set and recomputed all frequencies for that season. In total, we rejected $\approx 6\%$ of the observations acquired by the APT. The total number of observations given above is the number of observations that survived analyses of the individual observing seasons. These data are listed in Table 1. The full table is given on the VizieR website accompanying




Figure 1. The APT B and $B - V$ magnitudes for the γ Cas program for Seasons 1997–2019. Tick lines mark each season's end. The brightening and reddening segment after RJD 55,400 marks an outburst during 2010–2011. In the last season the $B - V$ color reddens because the Be star fades more in B than in V .

Table 1
APT Observations of γ Cas (Seasons 1997–2019)

Date (RJD)	Var B (mag)	Var V (mag)	Chk B (mag)	Chk V (mag)
50,718.6953	−4.306	−3.671	−0.809	−1.213
50,718.9258	−4.306	−3.672	−0.805	−1.219
50,720.7930	−4.299	−3.674	−0.812	−1.215
50,720.9180	−4.300	−3.668	99.999	−1.220

Note. A value of 99.999 indicates the differential magnitude had an uncertainty > 0.01 mag and was rejected.

(This table is available in its entirety in machine-readable form.)

this paper. The mean V and $(B - V)$ magnitudes in the Johnson system are $\langle V \rangle = 2.165$ and $\langle B - V \rangle = -0.083$.

4. Results

4.1. The Full Light Curve

The B differential magnitude data set is plotted in Figure 1. Besides short-term brightness variations, the plot exhibits a sinuous character over a range of ~ 0.05 mag in the 23 seasonal means. The bottom curve shows that the $B - V$ color index is reddened by 0.04 mag during this time, though not in strict correlation with B (or V) magnitudes. Because there is no other plausible source of surplus red continuum flux in the γ Cas system, we may assume that the variable reddening results from changes in disk extent and/or density. However, we also notice the unusual variations occurring between Seasons 2018 and 2019, when the B flux decreased and the V flux decreased too, but less so. It is possible that the disk has developed enough at that time that it has become opaque even in the blue. Then, assuming the inner disk edge occults part of the star, its enhanced optical thickness will dim the combined star/disk light even in the B passband.

4.2. Long Cycles

Table 2 lists by APT season the numbers of B and V observations and the lengths of the long cycles. These are taken

Table 2
Summary of APT Observations with Long Period and f_{82} Properties

Season	# Obsns. B/V	Long-cycle “Periods”	Long-cycle B/V Ampls.	f_{82} B/V Ampls.
1997	179/183	61	14.0/15.9	4.4/4.4
1998	206/209	65	6.8/7.6	5.9/6.1
1999	254/251	72	14.1/14.9	4.8/6.1
2000	290/290	91	14.8/17.6	3.4/6.1
2001	332/327	73	10.6/11.9	7.1/6.4
2002	300/300	80	16.4/21.0	7.6/4.7
2003	659/655	90:	19:/21: ³	6.3/7.2
2004	647/641	85	11.2/15.8	3.6/2.7
2005	287/275	66	6.0/4.9	5.3/2.6
2006	266/270	88	13.3/17.6	0.6/2.3
2007	254/248	88	11.2/13.9	0.8/2.0
2008	245/242	60	10.8/9.3	2.3/2.0
2009	192/188	70	10.2/9.3	0.7/1.5
2010	278/278	72	13.2/19.1	1.8/2.8
2011	326/318	73	17.7/20.1	1.5/2.0
2012	40/34	70	10.1/18.3	0.0/0.0
2013	93/93	1.4/0.0
2014	91/89	3.3/3.4
2015	132/126	3.4/2.9
2016	173/169	2.5/3.0
2017	87/88	2.2/0.0
2018	120/113	56	1.8/1.0	2.1/1.6
2019	103/101	73	1.6/1.9	1.5/2.4

Note. (1) For conciseness B and V properties are separated by a slash. (2) Full amplitudes are in millimagitudes; cycle lengths are in days. (3) Paper I showed damping/regrowth of cycle amplitude.

from Papers I and II and, from Season 2012 on, from new observations. As previously noted, the waveforms for the long cycles are not always sinusoidal. They can grow, damp out, or exhibit a net trend. At times they “morphed” to a new quasi-period within two weeks or less. Errors in the long-cycle lengths were estimated in Papers I and II to be ± 1 day for simple sinusoids to ± 2 days for damped cases. In contrast to those in Paper II, the full (peak-to-peak) amplitudes we list in Table 3 were computed by sinusoidal fits and without (sparse) summer observations. Thus the amplitudes here are not identical to those in the previous papers. The errors in cycle amplitudes are likewise dependent on the character of the variations and therefore also difficult to assign. We estimate them conservatively to be $\pm 15\%$ (see Figure 2). The full-amplitude detection threshold is about ~ 1 mmag.

From the seasonal history of the long cycles given in Figure 2 and the tables, their character appears chaotic, offering no obvious predictive power or memory of previous cycles. Following the cycles’ decline in 2012–2013, one sees that they may have recovered slightly in Seasons 2018–2019. This is at least consistent with the long-period, meandering character of the Transiting Exoplanet Survey Satellite³ (TESS) observations during this season, which further supports this general picture. Figure 3 shows that for cycle amplitudes larger than 10 mmag, the amplitude is larger in V than in B . There is no correlation of amplitude or amplitude color ratios with cycle length.

³ TESS (Ricker et al. 2015) was launched by NASA in 2018 to survey the sky with broadband optical photometry. The time cadence for γ Cas observations was 30 minutes.

4.3. Confirmation of f_{82} and Other Signals

4.3.1. Search Procedure

Our search procedure for coherent frequencies was first to run our Vaníček periodogram generator through a broad frequency range for each season (and each filter) and to tabulate the formal errors in the amplitudes of all significant peaks. This procedure worked well for f_{82} and for all but one of the other signals we may have found (see reference to 0.76 days^{−1} in Section 4.3.3).

We digress to point out that in their analysis of the TESS light curve of γ Cas during Sectors 17, 18, and 24, Labadie-Bartz et al. (2021, “LB21”) have discovered a low-amplitude group of nonradial pulsation (NRP) modes, which they designated as “group g_1 .” However, their amplitudes are too low to be detectable by the APT, and they flutter on an unknown timescale. The more stable and larger-amplitude f_{82} , first noted in the APT light curves of Papers I and II, occurs at the low-frequency edge of the g_1 group. Because these modes are so weak and well separated from aliases associated with the APT observing windows, we believe they are not directly related to f_{82} .

In Paper II we found that uncertainties in the full amplitudes for single seasons range from ± 1.0 to ± 1.5 mmag. After analyzing the individual seasons, we ran searches on three groups of seasonal data sets: early, middle, and late (see Table 3), according to the seasons added to Papers I and II and this work. The initial amplitude errors were formal ones, as propagated from the Vaníček analysis. The errors for multi-seasonal periodograms are lower than the single-season errors. The 23-season errors were calculated in the same manner.

To assess the effect of hypothetically fewer observations than were made, we split our database in two, each comprised of even- or odd-numbered points, and reran our 23-season analysis on the four discovered frequencies discussed below. As might be expected, the solutions for frequencies and amplitudes typically bracketed those computed from the full data set. The computed rms values based on the even-minus-odd observation differences varied between one and one and a half times the rms of the full-set solutions. In the “23 seasons” panel of Table 3 we have replaced the errors of the Vaníček solutions with those from the even/odd comparisons in cases where the former were larger. We also note that the amplitudes computed in the even/odd analyses fluctuated apparently randomly; there was no trend to smaller or larger values.

We repeated these trials four times by computing results for every fourth observation for these frequencies. About half these trials settled on the correct frequency, within the error windows given in Table 3. In the other cases a false peak (usually one of the adjacent annual aliases) was chosen, since the correct peak often did not stand out above them. With such a high failure rate as this, it was clear that we had reached the limit of our ability to detect astrophysical signals.

4.3.2. Discovery of Coherent Signals

The time history of f_{82} full amplitudes found in APT data is given in Figure 4 (values for 1997–2011 are from Paper II). Although this signal was strong in early seasons, the amplitudes decreased significantly after 2004–2005 to being barely visible until 2013 or 2014. From concurrent SMEI observations, Borre et al. (2020, “B20”) reported a similar decrease in the f_{82} amplitude. It appears to partially recover in

Table 3
Signal Frequencies (days^{-1}) and Full Amplitudes (mmag)

Multiseason	0.82	1.24	2.48	5.03
Paper I				
1997–2004 (<i>B</i>)	0.82245	1.24310	2.47944	5.02903
Full ampl.	5.92 ± 0.35	3.00 ± 0.37	1.53 ± 0.38	3.00 ± 0.38
1997–2004 (<i>V</i>)	0.82244	1.24304	2.47937	5.02328
Full ampl.	6.80 ± 0.39	3.30 ± 0.42	1.66 ± 0.42	3.31 ± 0.42
Paper II				
2005–2011 (<i>B</i>)	0.82929	1.24171	2.47975	5.03873
Full ampl.	2.28 ± 0.45	2.11 ± 0.45	2.23 ± 0.45	1.54 ± 0.46
2005–2011 (<i>V</i>)	0.81929	1.26146	2.47980	5.02477
Full ampl.	2.47 ± 0.45	2.36 ± 0.49	1.70 ± 0.49	1.34 ± 0.50
New data				
2012–2019 (<i>B</i>)	0.82394	1.24263	2.47946	5.03244
Full ampl.	2.32 ± 0.65	$1.58/2.37 \pm 0.65$	3.49 ± 0.64	2.25 ± 0.66
2012–2019 (<i>V</i>)	0.82762	1.25951	2.46901	5.02552
Full ampl.	2.46 ± 0.67	2.00 ± 0.68	2.44 ± 0.65	2.54 ± 0.66
23 seasons ⁽³⁾	0.82238(10)	1.2448(19)	2.481(11)	5.027(20)
<i>B</i>	3.51 ± 0.29	1.89 ± 0.29	2.42 ± 0.39	3.00 ± 0.38
<i>V</i>	3.58 ± 0.28	2.20 ± 0.29	2.15 ± 0.50	2.60 ± 0.41
f_{82} ephemeris from 23 seasons	T_0 (RJD)	P (days)	f_{82}	
<i>B</i>	51,086.602	1.215975	0.822385	
<i>V</i>	51,086.612	1.215987	0.822377	
Avg. T_0 , P , f	51,086.607(5)	1.21598(1)	0.82238(1)	

Notes. (1) The 1.21 day (f_{82}) ephemeris is $\phi = (T - T_0)/2\pi P$. (2) $\phi = 0.0$ refers to the “faint-star” phase. (3) The last digit in both the amplitude and frequency is not significant.




Figure 2. The seasonal APT history of long-cycle full amplitudes of γ Cas. Solid and dashed lines and symbols represent the filter. The seasons covered by Papers I and II are indicated. For most of the last several years the amplitudes have been too small to detect.




Figure 3. The correlation of *B* and *V* amplitudes in millimagnitudes from the long-cycle history in Figure 2. For amplitudes greater than 10 mmag the *V* amplitudes become significantly larger than the *B* amplitudes, demonstrating that the origin of these variations is the Be disk.

the following few years, but it is not visible in the later periodogram obtained during the TESS Sector 17–18 (2019–2020) observations (Nazé et al. 2020c, “N20c”). According to Figure 4, we cannot attest to the nonzero values for Seasons 2017–2019.

The lower panel of Table 3 gives our f_{82} ephemeris for all the data in both filters. The agreement between the f_{82} 23-seasons frequency for Seasons 1997–2019 and the three multiseason segments suggests that this frequency was coherent from when the APT monitoring began in Season 1997 through Season 2011 and probably into some late seasons. Our revised frequency reduces the discrepancy between the values reported

in Paper II and B20 ($0.82247 \text{ days}^{-1}$ versus $0.82215 \text{ days}^{-1}$) by one-third. These values now differ by close to 1.0 cycle over their timespan. It is likely that either they or we have miscounted by one cycle over the span of several thousand.

Other than 0.82 days^{-1} , we found multiseasonal signals near frequencies 1.24 days^{-1} , 2.48 days^{-1} , and 5.03 days^{-1} very similar, though not always identical, to results by B20, N20c, and LB21. Periodograms for frequencies surrounding these values are exhibited in Figure 5, and relevant parameters for them are listed in Table 3. The 2.48 days^{-1} signal seems to be a robust frequency for most, if not all, of the APT observing




Figure 4. The time history of the seasonal full amplitudes for the f_{82} signal for the B (X symbols) and V (squares) data sets. Errors for amplitudes of ± 1.2 mmag are averages of the seasonal values given in Paper II.




Figure 5. The B -filter full-program periodograms for the four frequencies found for most of the APT seasons covered in this paper. The comb astride the vertical line denoting these frequencies highlights the ± 1 annual aliasing pattern.

seasons. The 5.03 days^{-1} and 1.24 days^{-1} signals require additional notes.

N20c determined a peak value of 5.054 days^{-1} for $f_{5.03}$, which lies at our frequency error limit. Our periodogram for this signal shows evidence of stronger annual and daily aliases than the others. In marginal detection cases like this, these may be due to the frequency’s proximity to the diurnal harmonic at 5.0 days^{-1} .

The $f_{1.24}$ signal (B20’s $f_{1.25}$) was the most problematic to characterize. The filter-to-filter disparity in its amplitudes is large compared to results for the other signals, making our late seasons’ solutions for it less reliable. Indeed, this signal weakened in seasons after 2004. Moreover, we suspect its amplitude was generally highly variable. For example, N20c stated that in TESS Sectors 17–18 they found “no trace” of B20’s $f_{1.25}$. Yet, $f_{1.24}$ and $f_{2.48}$ signals were both present but variable at least for a few days during the beginning of Sector 18, as seen in their Figure 6 light curve. Finally, for this event or other times, we did not find that the APT or TESS

Table 4
Result Summary of Intensively Observed Nights

Season	# Nights	Freq.	Ampl.	Comments
2000	4	5.03	6	fits last 2 nights
2001	2	0.82	7	
2003	6	0.82	11	waveform change
2004	3	1.24(?), 0.82	12	evolving freqs.
2011	1	...	<2	flat over 5 hr
2016	2	2.48	18	

amplitudes of the two frequencies change together, as would be expected if the two were related through a harmonic resonance.

We searched our periodograms for additional signals (for example, the weak signal at 7.57 days^{-1} found by N20c in a TESS light curve) but detected none. In addition to the semi-stable frequencies just noted, we will cite a short-lived frequency in Section 4.3.3 and discuss an intermittent one in Section 4.6.

Before proceeding further, notice that there is no correlation between the amplitudes of the long cycles and the f_{82} signal. We note also that according to Table 1 and Pollmann et al. (2021), the disk of γ Cas has been building since 2000 through early 2021. In the middle of this interval, γ Cas underwent a Be outburst in 2010–2011 showing increased optical continuum and $H\alpha$ line brightening. As reported in Paper II, during the initial few weeks of this outburst, an accelerated brightening of the optical continuum, $B - V$ reddening, and $H\alpha$ emission occurred, which was accompanied by increased absorption in the soft X-ray region (SLM). In contrast, the near disappearance of f_{82} preceded the 2010 outburst by some five years and therefore was unrelated to that event. Thus, contrary to LB21, it is not clear that this event was associated with changes in strength of dominant NRP modes.

4.3.3. Rapid Amplitude Changes of Coherent Signals

The full amplitudes listed in Table 3 are values averaged over several seasons. From our short intensive-night campaigns (up to 7.5 hr per night), we found that the amplitudes can vary unexpectedly. An important result coming out of our intensive monitoring on short order is that the normally dominant signal at f_{82} was sometimes eclipsed by another, nominally secondary signal. Table 4 summarizes the results of several intensive mini-campaigns and the secondary frequencies (full amplitudes) they exhibited. Because we sometimes found temporarily dominant frequencies during these brief campaigns, their short-term behaviors suggest that their amplitudes vary much more frequently than we would infer from periodograms drawn from a full season or longer. Perhaps this “flutter” of signals is typical.

We discuss the results from our short dedicated-night campaigns season by season as follows:

Season 2000: This season included two pairs of intensive monitorings (each separated by two to three nights) spaced a month apart. Taking the second pair first, their variations could be fit with a signal of $f = 5.03 \text{ days}^{-1}$ and a mean full amplitude of 6 mmag. The behavior of the data in the first pair was decidedly different, exhibiting only small variations during their 4–4.5 hr coverages.

The B and V periodograms for the whole Season 2004 exhibited peaks of amplitude 4.8–5 mmag at 1.243 days^{-1} , as well as a strong transient signal (amplitudes 5–7 mmag) at a

frequency of 0.76 days^{-1} , in addition to the neighboring f_{82} . Though apparently real, this transient did not recur for any other season, and so we have not listed it in our tables as a multiseasonal signal.

Season 2001: The f_{82} full amplitude over two consecutive nights, 7 mmag, is typical for the season (Table 2).

Season 2003: The amplitudes of all three secondary frequencies were low or invisible during this season, signifying that the waveform changes discussed in Section 4.5.2 are unlikely to be due to intermode beating.

Season 2004: Light curves for a sequence of three consecutive nights were conspicuous with f_{82} appearing to beat with a sinusoid consistent with $\approx 1.2 \text{ days}^{-1}$. The combined full amplitude was large (12 mmag).

Season 2011: No variations were found over 5 hr.

Season 2016: The observations of these two consecutive nights were the only ones observed when the 2.48 days^{-1} signal was dominant. Its amplitude then is among the largest found during all our monitoring of γ Cas.

We temper these descriptions by noting that our fittings of large-amplitude sinusoids to data of only several hours of a few nights cannot be differentiated in general from beating by roughly similar modal amplitudes. The best single case for an amplitude waxing and waning within one to two weeks is discussed just below.

To summarize, from the well-observed nights referred to in Table 4, one sees frequent changes in amplitudes not only for f_{82} but also for the “secondary” signals near 1.24 days^{-1} , 2.48 days^{-1} , and 5.03 days^{-1} , all probably occurring on rapid timescales. It is not surprising to find in γ Cas what are evidently NRP modes excited in this frequency range, as it is a star situated at the hot edge of the β Cep domain. Although the f_{82} amplitude starts to decrease in 2005–2012, and is generally mimicked by $f_{1.24}$ and $f_{5.03}$, the amplitude of $f_{2.48}$ increases during later seasons (Table 3).

4.4. Searches for Other Frequencies

High frequencies ($> 8 \text{ days}^{-1}$) are of interest to this study because they are potentially identified with NRP p-modes and therefore possibly cause msf in line profiles (Nazé et al. 2020b, “N20b”). However, the quality of our APT periodograms deteriorates above $8\text{--}10 \text{ days}^{-1}$ and is meaningless beyond it. However, N20c’s published TESS periodograms for Sectors 17 and 18 exhibit no signal in the range above 8 days^{-1} out to 20 days^{-1} (just as LB21’s periodograms show no high-frequency signal down to $< 0.1 \text{ mmag}$, not only for this time period but also during Sector 24).

4.5. Changes in f_{82} Waveform

4.5.1. Rationale for Analysis

In Paper I we discovered an unusual skewness in the mean waveform of the f_{82} signal taken from the first several APT seasons. For convenience we characterized departures from a sinusoid by parameters e and ω taken from the familiar Lehmann-Filhes equation for orbital solutions of radial velocity variations. Here e and ω are a fake eccentricity and longitude of periastron, respectively. Parameter e represents the waveform’s pointiness while ω quantifies its skewness. Values in the fourth quadrant signify a depressed positive-phase wing; the first quadrant gives the opposite skewness. In our implementation of past and current work, we used a

generalized least-squares algorithm by Markwardt (2011) adapted for our computations. For the first eight seasons (Paper I), the resulting means, averaged over B - and V -filter data sets, were $e = 0.35$ and $\omega = +285^\circ$. In Paper II we examined data for six dedicated and consecutive nights of observations in Season 2003 and found for the V filter that e had increased to 0.51 ± 0.05 and the skewness had reversed signs to $\omega = +25^\circ \pm 6^\circ$. As noted then, nearly identical values and errors were found in our data by Dr. Fekel using an independent algorithm. The departures from a sinusoid are a remarkable result and thus require confirmation. We note for completeness that the periodogram for these six nights’ data exhibits a faint second-harmonic feature.

To check the statistics in a different way, we conducted an experiment adopting a simple Monte Carlo strategy for the V -band data set of another observing season, 2001, and compared errors derived for e and ω from fake data sets using the 2001 observation times and photometric errors. We then compared them with results from a direct analysis of Season 2003 data. Note that an assessment of fluctuations of both seasons’ target and check-minus-comparison star data (with outliers removed) revealed no significant departures from Gaussian distributions.

Our experiment began by phase-folding the Season 2001 V -band data to the f_{82} ephemeris. The best (though mediocre) sinusoidal fit was computed for these initial data, and this sinusoid was subtracted, resulting in an initial data fluctuation array. We then conducted mock data simulations for 11 independent trials by shifting the fluctuation array by phase increments of $N \times 28$ points according to each point’s former position. Here $N = 0, 1, 2, \dots, 10$ corresponds to the trial number and 28 is an arbitrary value that brings each point to a different observation time and night. For each trial, fluxes of the so-shifted fluctuation array were added back to the initial fitted sinusoid, and e and ω were solved for again. Finally, their means and rms errors from these trials were computed and compared with the original e and ω computed for Season 2001. The mean values turned out to be nearly coincident with the solution for the original Season 2001 data, namely means of $e = 0.16$ and $\omega = 310^\circ$ and rms errors of $\sigma_e = \pm 0.077$ and $\sigma_\omega = \pm 8^\circ$.

We repeated this exercise by sampling only every other observation. The new rms values were $\sigma_e = \pm 0.102$ and $\sigma_\omega = \pm 12^\circ$ and thus scaled approximately with the inverse square root of the number of points. Applying the same scaling for like-quality and increased observations to Season 2003 (Table 2), we were able to predict errors of $\sigma_e = \pm 0.055$ and $\sigma_\omega = \pm 6^\circ$. These error estimates are nearly the same as Paper II’s results from a direct analysis of Season 2003 (viz., ± 0.05 and $\pm 6^\circ$). We will now use this result in the following analysis of data subsets of this season.

4.5.2. Season 2003 Waveform Changes

We proceeded to analyze the waveform for B - and V -band observations of our six intensively monitored nights from Season 2003 (RJD 52,962–52,967). We will then contrast it to observations taken from the rest of the season. In comparison to Paper II’s analysis, we incorporated differences in outliers comprising the seasonal data set and used different procedures for prewhitening of the long period in this analysis. We remind the reader that we found the amplitudes from the secondary signals to be low or absent in this season.

Table 5
Lehmann–Filhes Waveform Parameters, Season 2003

Filter	Halves 1 + 2	(All but 6 Nts.)	Intensive	(Only 6 Nts.)		
	Ampl.	e	ω	Ampl.	e	ω
<i>B</i>	5.2	0.41	348°	12.1	0.50	+20°
<i>V</i>	4.2	0.41	307°	10.2	0.47	+19°
	Half 1			Half 2		
	Ampl.	e	ω	Ampl.	e	ω
<i>B</i>	5.5	0.34	350°	5.7	0.43	340°
<i>V</i>	5.1	0.39	340°	5.1	0.41	319°

Note. The set of six intensively monitored nights occurred between two nearly equal time segments during this observing season.

To examine the waveform differences during this season, we used the fact that the six intensively monitored nights occurred in the middle of the observing season. We divided the data sets for the nonintensively observed nights into two halves and compared the resulting waveforms of all four groups—viz., the first half, the second half, both halves (all nonintensive nights), and the six intensive nights. The e and ω parameter determinations for these groups are given in Table 5. The most obvious result is that the full amplitude doubled from 5 to 6 mmag to ~ 11 mmag, and then decreased to its former value. Also, the “eccentricity” for the six nights increased to 0.50 and 0.47 for *B* and *V*, respectively. This increase in e , +0.16, is more than double the predicted σ_e of ± 0.055 , according to our control results for Season 2001 (when scaled for numbers of data points).

These differences are visible in the phase-folded plot in Figure 6. Here the Lehmann–Filhes fit to the six-night, *B*-filter data points is displayed as a solid line. We can contrast it with the Lehmann–Filhes fit to the data for all other nights in the season (dashed line) and also with the best, though mediocre, sinusoidal fit to the data for these other nights (dotted–dotted–dashed line). The differences between the values of e and ω and the published values in Paper II are comparable to the error estimates found in our control: $e = 0.50$ here versus 0.55 in Paper II, and $\omega = +19^\circ$ here versus 25° there. Also, just like the eccentricity, the skewness subsequently reverted to its typical fourth-quadrant value.

4.6. The UV Continuum Dips

Although the two IUE UV light curves of γ Cas lasted longer, the GHRs series of 1996 March 14–15 is unmatched in its precision. In Figure 7 we have exploited this fact and the f_{82} ephemeris to represent this signal as a 6 mmag sinusoid (dotted line) against the GHRs data (first full curve). The error in the phase positioning of the sinusoid is ± 0.01 cycles. We have subtracted the sinusoid from it to show how it would look if f_{82} were not present (second full curve). This would be appropriate, for example, if f_{82} is a very low frequency, “classical” NRP g-mode, for which geometric distortion of the star would dominate flux variations. The amplitude of the variations would then be wavelength-independent. If instead, as argued by SRH, the absorptions arise from a cool intervening cloud, then wavelength dependence grows in the far-UV, and the GHRs light curve will approximate the lower curve. In any case, the lower curve gives an approximation of a sinusoid, for which least-squares fitting gives a dip separation of 8.90 ± 0.02 hr (2.70 ± 0.01 days $^{-1}$). However, it is easy to show from the 1982 and 1996 IUE records, and even from the GHRs curve,




Figure 6. The waveform determined by least-squares fit to the Lehmann–Filhes solution for the *B*-filter, f_{82} phase-folded data for six intensively monitored nights in Season 2003 (heavy curve and dots). Note the departure of the data and this curve from the sinusoidal fit (dotted–dotted–dashed) and the Lehmann–Filhes fit (dashed) for the nonintensive nights: the six-night solution has a larger amplitude and exhibits a “pointy/bowed” form and skewness with a steep negative-phase wing.




Figure 7. The GHRs UV continuum light curve of 1996 March 15 (upper full curve). The top line (dotted sinusoid) is the f_{82} signal from its ephemeris of Table 3. The second and third curves are the original GHRs curve with the top sinusoid subtracted by a factor of 1 and 2.5, respectively. One or the other of the lower curves shows how the undistorted UVC should appear if no f_{82} signal existed.

that these variations are not part of a true sinusoidal signal. Even so, S19 pointed out that the 43 hr long IUE sequence suggests the presence of a third dip that seems to be the recurrence of the first dip from one rotation cycle earlier.

One might ask whether the 8.9 hr separated dips are coherent enough through the years to be detected in the APT data, since the program started just over one year after the GHRs campaign. Yet even if they are present, they are not continuously repeating sinusoidal signals. An intermittently occurring dip will cause the periodogram to show a more complicated beating structure. Fortunately, because we now know where to isolate a narrow search range, we can search a periodogram for a peak that emerges at a predicted frequency of 2.70 days $^{-1}$. We first searched for signals in the *B* and *V* periodograms in the 23-season composite and found none. However, we did find a possible weak peak at 2.695 days $^{-1}$ in




Figure 8. Periodograms for B (dashed line) and V (solid line) covering a weak signal at the expected frequency 2.695 days^{-1} . Panel (a): Results for Seasons 1998–2003. The horizontal line and markers indicate the feature’s annual sidelobes. Panel (b): The same feature for just Season 1997. Panel (c): The simulation of the GHRS signal applied to Season 1997 (see text). For panels (b) and (c) we extend the frequency range to $2.5\text{--}2.82 \text{ days}^{-1}$ to show the noise level. (The slight positional differences of the signal are not significant.) The dotted line in panel (b) is the extension of the signal arrow in panel (a). This position is almost identical to that of the dotted-dashed line (panel (c)).

the 1998–2004 multiseason periodograms, which are shown in Figure 8(a). Their annual sidelobes are also visible. To see if this candidate signal also exists for Season 1997, the season closest to the UV 1996 campaigns, we computed the periodograms for this season and display them in Figure 8(b) to show that they have the same peak. Note that although the Season 1997 noise is twice as large as the Season 1998–2004 noise, the signal is three times as strong. From these results, the putative signal is weakening and becomes invisible in late APT seasons. Although there is no immediate way of checking, it is possible that in recent years the UV dips have no longer been present.

To verify our Season 1997 detection, we have constructed artificial data sets by estimating the egress wing to the pre-dip level and thus completing the expected UVC curve out to 1.216 days from observation start (see dashed line in Figure 7). We then repeated and concatenated the curve 115 times until it covered the full span of this season. Since the S/N of the GHRS greatly exceeded the APT’s, we then added Gaussian noise to simulate the APT’s data fluctuations. Finally, we sampled the noisy, season-long curve at the actual observation times. We repeated the procedure for various assumed noise levels. The result was a series of mock light curves for various noise levels that retain the observing window gaps.

Figure 8(c) shows the resulting periodogram for a mock S/N of 250 per observation, which matches the measured APT rms. The position of the generated signal, at 2.697 days^{-1} , is already “locked in” to the 8.90 hr we measured earlier, so its position is a given. Similarly, the S/N of the peak height to the surrounding noise level is not unexpected because the estimated noise level of the observations was determined from the scatter of comparison-check star observations (see Section 3). However, the near agreement of the simulated and observed peak heights in Figures 8(b) and (c) reassures us that the signal is stellar. Because the strengths of the dips are

already known to vary over time (e.g., the dips were stronger in 1982), we consider this amplitude agreement fortuitous.

5. Discussion

5.1. High-frequency Pulsations and Msf

High-frequency modes are now known to be active in Be stars, e.g., in π Aqr, a γ Cas analog, which exhibits one or more tesseral modes (N20b). The amplitudes of these features traveling through line profiles amount to several percent in line profiles (1–2.5 mmag in the optical light curve). In N20b’s Figure 4 one sees that the spacings of NRP-induced msf in optical spectra of π Aqr, undoubtedly due to high-frequency p-modes, are rather uniformly spaced in time. Thus, there are more differences than similarities in the msf of this star’s spectrum compared to γ Cas.

In view of the results on π Aqr, we now revisit the Yang et al. (1988) and our own reports of msf in optical and UV spectral profiles of γ Cas. In this context, L20 doubted our interpretation of these features as clouds forced into corotation over transient magnetic centers on the star (SR99). Therefore, we now address arguments for an NRP interpretation for msf.

The most precise record of msf in the spectrum of γ Cas is the 21.5 hr time series of GHRS UV spectra. In this series, a raft of msf associated with many lines is ubiquitous. The features travel across line profiles at a rate of $+95 \pm 5 \text{ km s}^{-1} \text{ days}^{-1}$ and reoccur at unpredictable intervals averaging very roughly 2 hr (12 days^{-1}) during this time series. Importantly, they wax and wane in visibility in about 1.5 hr. In a separate study Smith (1995) discussed 109 high-dispersion difference spectra of the He I $\lambda 6678$ profile during five nights in 1993; each monitoring interval was 3–6 hr. The difference spectra revealed msf striation patterns that reoccurred at erratic intervals on most nights and lasted no more than 2–3 hr. The acceleration rate of the features was $+92 \pm 10 \text{ km s}^{-1} \text{ days}^{-1}$, in good agreement with the later UV results. Similarly, the unpredictable appearances and short lifetimes of these events render any attempt to measure their recurrences all but meaningless. The cyclical intervals between these patterns averaged 2–2.5 hr in the 1993 monitoring and 1.5–2 for the 1996 UV monitoring. For the following discussion we note that the observed amplitudes of the msf were about 0.4% for the optical He I line and (depending on the line’s excitation) 0.3%–0.6% for the UV, i.e., the msf amplitudes are similar in the two wavelength regimes.

With this description, we discuss why high-frequency NRPs are not the best explanation for the msf in γ Cas:

1. The absorption features are noncoherent. Also, unlike NRP bumps in line profiles, they are not necessarily most prominent in the middle of their lifetimes.
2. To match roughly the irregular spacings of the msf in the GHRS data set, an NRP p-mode would have a frequency of $9\text{--}12 \text{ days}^{-1}$. The reported signal at 7.57 days^{-1} (N20c; LB21) is too low to meet this criterion.
3. Consider that the ratio of line profile msf semi-amplitudes to photometric msf semi-amplitudes in π Aqr is 2.5% to 1 mmag. These variations are due to NRP (N20b). Because the amplitudes of msf in γ Cas spectra are five times smaller, the photometric amplitudes in γ Cas periodograms should be 0.2 mmag if they too are caused by NRP. Since the TESS periodogram of γ Cas rules out photometric amplitudes down to less than

0.1 mmag at high frequencies, the line profile msf of γ Cas are not likely to be caused by NRP.

4. Within measurement errors, the amplitudes of msf are the same for optical and UV spectra. This is an important point because the restoring force of high-frequency pulsations is due to pressure imbalances (from their high frequencies they are likely to be p-modes). Model atmosphere simulations indicate that temperature variations from pulsations represented by spherical harmonics cause flux variations in early-type B stars that are approximately two and a half times as large as those in the optical, not equal to them as observed.

Of these arguments, the third one is probably the most powerful. However, the stipulation should be made that the amplitude of the line profile and photometric msf reflected behaviors at different times. The fourth argument relies on the wavelength dependence of amplitudes in the UV being large compared to that in the optical for high-degree p-modes, as is true for low-degree ones. At least for rapidly rotating, early Be stars this is relatively unexplored territory, and further exploration is necessary. To date, the amplitude behavior with wavelength seems to have been investigated theoretically only up to intermediate-degree ($l=3-4$), classical g- and p-modes (e.g., Pigulski et al. 2017).

We remark further that if UV-absorbing structures are suspended over different stellar latitudes, their signatures will exhibit more than one acceleration across the line profiles.

5.2. The Nature of the 0.82 days^{-1} Frequency

As part of our initial justification for assigning f_{82} to rotational modulation, Paper I argued that the alternative, classical NRP modes were not likely to be excited in rapidly rotating early-type Be stars, whereas this frequency is quite consistent with the rotational modulation of a surface inhomogeneity. As described next, this was likely to be a premature conclusion. In the meantime, various spectroscopic campaigns as well as a flood of results and satellite photometric surveys have shown that NRPs are endemic to Be stars, including early-type and rapidly rotating stars.

NRP was given a major boost from the study of spectral line profile variations for a small but representative population of other early-type classical Be stars (e.g., Rivinius et al. 2003). More recently, the discovery of groups of frequencies sometimes close to the rotation frequency, Ω_R , has presented evidence that nearly all of them must be due to NRP (Baade et al. 2016). Even so, some investigators (e.g., Balona & Engelbrecht 1986; Balona 1995) have argued that one of these frequencies, particularly if it is visible at intermittent intervals, can be caused by the rotational modulation of a starspot.

Satellite photometry has demonstrated that NRP modes are endemic to early-type Be stars. Recent photometric satellite surveys (e.g., Balona et al. 2015; Semaan et al. 2013, 2018; Saio et al. 2017; Labadie-Bartz et al. 2020; Balona & Ozuyar 2020a, 2020b, “BO20b”) have disclosed that, for most stars exciting low-frequency signals near their rotation rates (Ω_R), these signals are members of clusters of frequencies, wherein only one at most can be rotational.

Yet, one might ask whether there exist rapidly rotating, early-type Be stars with single isolated frequencies near Ω_R . L. A. Balona (2020, private communication) reports that in the larger TESS survey of BO20b, five O9–B2 stars exhibit

apparent isolated, coherent modes at frequencies that arguably coincide with the rotation frequency. Therefore, because such signals that meet our conditions do exist in a few early-type Be stars, it is now apparent we can no longer hold, as in our previous papers, that the isolated signal at f_{82} , though arguably close to Ω_R , is unique to γ Cas.

Before interpreting the f_{82} frequency as arising from a long-period g-mode, one should consider a potential obstacle: an oscillation observed near f_{rot} in the inertial frame will have a frequency of nearly zero in the corotating frame, i.e., the period will be very long in the (physically important) reference frame. Its corresponding peak in the periodogram would blend with its high-order neighbors, causing a broad peak, which is not observed. A better identification would be of an r-mode, which is an essentially horizontal vortical pattern at the surface excited by Coriolis force imbalances (or the κ mechanism; Saio et al. 2017), which we turn to next.

Several years ago Walker et al. (2005) reported the excitation of a thicket of low frequencies ($\lesssim 0.005 \text{ mHz}$) in the rapidly rotating B5e star HD 163868. Unlike several other modes of frequencies 0.02 mHz or higher, which can be ascribed to p- or g-modes, this low-frequency cluster is close to a multiple of the rotational frequency (i.e., $0.90-0.95 m\Omega$, where perhaps $m=1$). According to Saio (2013, 2018), these frequencies are probably signatures of odd, low azimuthal order r-modes. The circulation of surface particles participating in r-modes is associated with large polar-direct velocity components at midlatitude. This renders the discovery of r-modes particularly advantageous in stars observed at intermediate inclination like HD 163868.

Since γ Cas is likewise observed at an intermediate inclination, its f_{82} signal may likewise be due to a low-degree r-mode. Notably, r-modes are predicted to occur as clusters of frequencies, often adjacent to a dominant one (e.g., Walker et al. 2005; Saio 2018). From previous and present APT results, it appears to us as if only an isolated peak is visible near f_{82} . If f_{82} is indeed an r-mode, other associated r-modes may be present with amplitudes too small to be detected by the APT. Thus, LB21’s detection of multiple low-amplitude g_1 -modes in their TESS light curve appears to be consistent after all with f_{82} being the dominant mode of an r-mode complex, at least when it was visible in the early years of the APT program. In addition to γ Cas and HD 163868, a few rapidly rotating B stars in the cluster NGC 3766 may well excite both g- and r-modes (Saio et al. 2017).

5.3. Extra-APT Contributions

As few as the UV satellite monitorings of γ Cas are, combined with the revised ephemeris of Table 3, they permit a reinvestigation and an entry of new evidence as to the origin of the f_{82} signal.

We measured the times corresponding to passages of the centroids of the first dip in the IUE 1982, IUE 1986, and GHRS light curves (Figure 7). These occur, respectively, at RJD 44,997.48, 50,101.57, and 50,157.70. According to our Table 3 ephemeris, these times correspond to faint-star phases of 0.41, 0.92, and 0.08, respectively. We estimated errors on the 1996 IUE and GHRS phases as ± 0.02 and ± 0.03 . These values are dominated by errors in our adopted frequency (Table 3). For the more important phase difference error between the two 1996 first-dip centroids (the time interval being 57 days), the error in the frequency is negligible. The error in the 1996 IUE

feature relative to the GHRS feature is dominated by centroid measurement and is ± 0.02 . The errors for the 1982 IUE dip are larger, ± 0.05 , because the frequency and centroid-finding errors must be folded together. Similarly, the phases for the centroids of the second UV dip are 0.71, 0.22, and 0.39. We estimated phase errors in the GHRS second dip to be dominated by uncertainty in frequency, whereas errors in the IUE second-dip centroids are about 50% larger because their profiles are not as well defined. If we had instead adopted the B20 frequency and our T^* from Table 3, the phases would be different, but the net result would be much the same. All told, the far-UV features are not phase-locked with the optical ephemeris. The phasing mismatch occurs even over the small interval (46.16 cycles) in the ≈ 57 day interval between the 1996 IUE and GHRS observations. Assuming a cycle miscount of 0 or 1, the percentage mismatch would be either 0.4% or 2.5%, respectively, which, though small, introduces a phase slippage.

The resolution of this slippage starts with the fact that only one frequency can be rotational. We believe it is unrealistic that, over the 57 day interval between the 1996 IUE and GHRS observations, a surface differential rotation rate of order 1% or more occurs. Therefore, we reject this possibility. It follows that we prefer to adopt the alternative: f_{82} is not a rotational signal. Yet, our preference does not prove the case. In fact, given the isolation of the signal at low frequency and its sometime nonsinusoidal waveform, the identification of f_{82} with NRP is not straightforward. In any case, in view of the arguments put forth regarding the UV color changes of the dips (SRH) and the correlation of appearances with changes in UV spectral lines and hard X-ray flux (SLM16), we can see no reason to reject the corotating picture. This does not mean that the rotation frequency has been found. Given the physical parameters of the star and the likely reoccurrence of the first dip in the 1982 IUE light curve, it is probably near 0.8 days^{-1} .

6. Summary and Conclusions

We summarize the main points of this work as follows:

1. The discovery of coherent signals with frequencies at 1.24, 2.48, and 5.03 days^{-1} by B20 and N20c has been confirmed by an independent data set (APT).
2. We agree with previous authors that these signals are NRP modes (p-modes or low-degree g-modes). These modes should not be confused with the stochastic low-frequency ($\sim 0.1\text{--}2 \text{ days}^{-1}$) variability discovered in a variety of OB stars in TESS data (e.g., Bowman et al. 2020). For early B stars the amplitude of such “white noise” is generally only $\sim 0.1 \text{ mmag}$ in B III–V stars and thus is well below the detection limit of the APT. This variability is thought to be excited by turbulence generated within the Fe-opacity convective zone (e.g., Cantiello et al. 2021).
3. The $0.82238 \text{ days}^{-1}$ frequency was stable in the *B* and *V* filters from 1997 through 2011, although the amplitude varied. Since then, the signal has faded and then shown a weak recurrence (2014–2016). We cannot verify that it was active during 2017–2019.
4. NRP amplitudes can at times wax and wane rapidly. Such activity is also displayed in N20c’s TESS dynamic periodogram and Figure 9.

5. We believe the 0.82 days^{-1} signal is an excited NRP mode of a still undetermined type. However, its occasional tendency to modify its waveform complicates a physical description of its origin. Also, the isolated position of a detectable signal near Ω_R (but how near?) may not be unique among early Be stars (BO20b), but it is not the norm either. Otherwise, given the excitation of these modes, γ Cas has begun to resemble other Be stars at the periphery of the β Cep domain.
6. Our preference for attributing f_{82} to NRP rather than to rotational modulation was facilitated by using UV light curves. These indicate phase shifts from our f_{82} ephemeris. We identify the pair of dips from UV photometry as being likely due to rotation. However, because we do not know the putative stellar longitude separations of the absorbing structures, we cannot determine the exact value of Ω_R .
7. In periodograms of early APT season data we found a signal at 2.70 days^{-1} that corresponds to the time separation of the two-dip pattern observed in the three UV campaigns of 1982 and 1996. However, periodograms of (most) later APT seasons and recent satellite data sets do not exhibit this signal. We conclude that it has diminished and may no longer be visible in the UV or optical.
8. TESS data offer no evidence that high-frequency NRP modes produce the msf in optical and UV spectra of γ Cas. Moreover, the msf are chaotic, exhibit much larger amplitudes than the APT and TESS detection thresholds, and do not show an expected increase in amplitude from the optical to the far-UV.
9. The so-called long cycles abruptly faded to invisibility just after the era covered by Paper II and shortly after the 2010 outburst. (A possible recovery in two recent seasons might have occurred (Figure 2) but is too weak to be reliable.) We speculate that the continued buildup of the inner disk, according to APT photometry and especially the increased He I $\lambda 6678$ emission (Pollmann et al. 2014), is caused by an increased density there that overwhelms a fragile disk dynamo mechanism.
10. The correlation of APT and X-ray long cycles argues that the Be disk mediates the production of hard X-rays on the star. Optical variations found by the APT have therefore been important in framing the magnetic interaction hypothesis.

In criticizing the star–disk magnetic interaction hypothesis, L20 and B20 overlooked some key points. The existence of the f_{82} signal is largely irrelevant to the production of hard X-ray flux in this picture.⁴ The spectral msf are supportive of, though not essential to, the basic picture unless they can be identified with large X-ray “flares.” However, if instead they turn out to be due to high-frequency NRPs after all, the case for suspended cloudlets would disappear. The optical/X-ray long-cycle connection is important to the picture—the previous points are not required. The high densities associated with the flares strongly suggest a photospheric origin, to say nothing of the correlation of hard X-ray fluxes with photospheric UV line strengths (SR99). Therefore, any

⁴ The rotation rate comes into the discussion of the details of corotating clouds, and then only secondarily in estimating their elevations. As such the error in equating f_{82} to Ω_R cannot be large.

rapid, aperiodic events, e.g., those caused by emerging magnetic structures, should be examined as aiding in the understanding of the X-ray formation process.

This study would not have been possible without the support of Lou Boyd, who has conscientiously managed the automatic telescopes at Fairborn Observatory. We are indebted to Drs. Yael Naze, Gregor Rauw, and Andrzej Pigulski and Mr. Piotr Kolaczek-Szymanski for sending us satellite light-curve extractions of γ Cas and for many suggestions that improved the manuscript. We are also grateful to Drs. Raimundo Oliveira de Lopes and Christian Motch for comments on the draft. It is a pleasure to acknowledge an anonymous referee's calling our attention to an important error in an earlier version of the manuscript and the referee's many insightful suggestions. G.W. H. acknowledges long-term support from NASA, the National Science Foundation, Tennessee State University, and the State of Tennessee through its Centers of Excellence Program.

ORCID iDs

M. A. Smith  <https://orcid.org/0000-0003-4240-4794>

G. W. Henry  <https://orcid.org/0000-0003-4155-8513>

References

- Baade, D., Rivinius, Th., Pigulski, A., et al. 2016, *A&A*, **588**, 56B
- Balona, L. A., 1995, *MNRAS*, **277**, 1547B
- Balona, L. A., Baran, A. S., Dasyńska-Daszkiewicz, J., et al. 2015, *MNRAS*, **451**, 1445B
- Balona, L. B., & Engelbrecht, C. A. 1986, *MNRAS*, **219**, 131B
- Balona, L. B., & Ozuyar, D. 2020a, *MNRAS*, **493**, 2528
- Balona, L. B., & Ozuyar, D. 2020b, arXiv:2008.06288v1
- Borre, C. C., Baade, D., Pigulski, A., et al. 2020, *A&A*, **635**, A140
- Bowman, D. M., Burssens, S., Simón-Díaz, S., et al. 2020, *A&A*, **640**, A36
- Cantiello, M., & Braithwaite, J. 2011, *A&A*, **540**, A140
- Cantiello, M., Lecoanet, D., & Jermyn, A. S. 2021, arXiv:2102.05670v2
- Doazan, V., Franco, M., Sedmak, G., et al. 1983, *A&A*, **128**, 171
- Eaton, J. A., Henry, G. W., & Fekel, F. C. 2003, *ASL*, **288**, 189E
- Gies, D. R., Bagnuolo, W. G., Ferrara, E. C., et al. 1998, *ApJ*, **493**, 440G
- Hamaguchi, K., Oskina, L., Russell, C., et al. 2016, *ApJ*, **832**, 140H
- Harmanec, P. 2002, in ASP Conf. Ser. 279, Exotic Stars as Challenges to Evolution, ed. C. Tout & W. Van Hamme (San Francisco, CA: ASP), 221
- Harmanec, P., Habuda, P., Stefl, S., et al. 2000, *A&A*, **364**, L85
- HEASARC Rosat_Data_Archive 2020, <https://heasarc.gsfc.nasa.gov/docs/cgro/db-perl/W3Browse/w3table.pl>
- Henry, G. W. 1995a, in ASP Conf. Ser. 79, Robotic Telescopes. Current Capabilities, Present Developments, and Future Prospects for Automated Astronomy, ed. G. W. Henry & J. A. Eaton (San Francisco, CA: ASP), 37
- Henry, G. W. 1995b, in ASP Conf. Ser. 79, Robotic Telescopes: Current Capabilities, Present Developments and Future Prospects for Automated Astronomy, ed. G. W. Henry & J. A. Eaton (San Francisco, CA: ASP), 44H
- Henry, G. W. 1999, *PASP*, **111**, 845
- Henry, G. W., & Smith, M. A. 2012, *ApJ*, **760**, 10
- Labadie-Bartz, J., Baade, D., Carciofi, A. C., et al. 2021, *MNRAS*, **502**, 242
- Labadie-Bartz, J., Carciofi, A. C., Henrique de Amorim, T., et al. 2020, arXiv:2010.13905v1
- Langer, N., Baade, D., Bodensteiner, J., et al. 2020, *A&A*, **633**, A40
- Mamajek, E. 2017, *JDSO*, **13**, 264
- Markwardt, C. 2011, Markwardt IDL Library, <https://pages.physics.wisc.edu/~craigm/idl/idl.html>
- Mason, K. O., White, N. E., & Sanford, P. W. 1976, *Natur*, **260**, 690
- Motch, C., Lopes de Oliveira, R., & Smith, M. A. 2015, *ApJ*, **806**, 177
- Murakami, T., Koyama, K., Inoue, H., et al. 1986, *ApJL*, **310**, L31
- Nazé, Y., Motch, C., Rauw, G., et al. 2020a, *MNRAS*, **493**, 2511
- Nazé, Y., Pigulski, A., Rauw, G., et al. 2020b, *MNRAS*, **494**, 958
- Nazé, Y., Rauw, G., & Pigulski, A. 2020c, *MNRAS*, **498**, 3171
- Nemravová, J., Harmanec, P., Koubska, P., et al. 2012, *A&A*, **537**, A59
- Peters, G. J., Gies, D. R., Grundstrom, E. D., et al. 2008, *ApJ*, **686**, 1280
- Peters, G. J., Pewett, T. D., Gies, D. R., et al. 2013, *ApJ*, **765**, 2P
- Pigulski, A., Baran, A., Bzuowski, M., et al. 2017, in Proc. Polish Astron. Soc. 5, ed. K. Zwintz & E. Poretti, 76
- Pollmann, E., Vollmann, W., & Henry, G. W. 2014, *IAUIB*, **6109**, 1
- Pollmann, E. 2021, *JAVSO*, **49**, 1
- Postnov, K., Oskina, L., & Torrejón, J. M. 2017, arXiv:1017.00336v1
- Ricker, G. R., Winn, J. N., Vanderspek, R., et al. 2015, *JATIS*, **1**, 014003
- Rivinius, Th., Baade, D., & Stefl, S. 2003, *A&A*, **411**, 229
- Robinson, R. D., & Smith, M. A. 2000, *ApJ*, **540**, 474
- Robinson, R. D., Smith, M. A., & Henry, G. 2002, *ApJ*, **575**, 435
- Saio, H. 2013, in Seismology for Studies of Stellar Rotation, Vol. 835 ed. K. Belkacem, 3M
- Saio, H. 2018, in Proc. Physics of Oscillating Stars Conf. arXiv:1812.01253
- Saio, H., Ekström, S., Mowlavi, N., et al. 2017, *MNRAS*, **467**, 3864
- Semaan, T., Hubert, A. M., Zorec, J., et al. 2013, *A&A*, **551**, 130
- Semaan, T., Hubert, A. M., Zorec, J., et al. 2018, *A&A*, **617**, A70
- Smith, M. A. 1995, *ApJ*, **442**, 812
- Smith, M. A. 2019, *PASP*, **131**, 044201
- Smith, M. A., Henry, G. W., & Vishniac, E. 2006, *ApJ*, **647**, 1375
- Smith, M. A., Lopes de Oliveira, R., Motch, C., et al. 2012, *A&A*, **540**, A53
- Smith, M. A., Oliveira, R., & Motch, C. 2016, *ASpR*, **58**, 782
- Smith, M. A., & Robinson, R. D. 1999, *ApJ*, **517**, 866
- Smith, M. A., & Robinson, R. D. 2003, in ASP Conf. Ser. 292, Interplay of Periodic, Cyclic, Stochastic Variability, ed. C. Sterken (San Francisco, CA: ASP), 263
- Smith, M. A., Robinson, R. D., & Corbet, R. H. 1998a, *ApJ*, **503**, 877
- Smith, M. A., Robinson, R. D., & Hatzes, A. P. 1998b, *ApJ*, **508**, 945
- Vančec, P. 1971, *Ap&SS*, **12**, 10
- Walker, G. A., Kuschnig, R., Matthews, J. M., et al. 2005, *ApJL*, **635**, L77
- Wang, L., Gies, D. R., & Peters, G. J. 2017, *ApJ*, **843**, 60
- Wang, L., Gies, D. R., Peters, G. J., et al. 2021, *AJ*, **161**, 248
- White, N. E., Swank, J. H., & Holt, S. S. 1982, *ApJ*, **263**, 277
- Yang, S., Ninkov, Z., & Walker, G. A. H. 1988, *PASP*, **100**, 233