

## MEASUREMENT OF THE SPIN-ORBIT ALIGNMENT IN THE EXOPLANETARY SYSTEM HD 189733<sup>1</sup>

JOSHUA N. WINN,<sup>2</sup> JOHN ASHER JOHNSON,<sup>3</sup> GEOFFREY W. MARCY,<sup>3</sup> R. PAUL BUTLER,<sup>4</sup> STEVEN S. VOGT,<sup>5</sup>  
GREGORY W. HENRY,<sup>6</sup> ANNA ROUSSANOVA,<sup>2</sup> MATTHEW J. HOLMAN,<sup>7</sup> KEIGO ENYA,<sup>8</sup> NORIO NARITA,<sup>9</sup>  
YASUSHI SUTO,<sup>9</sup> AND EDWIN L. TURNER<sup>10</sup>

Received 2006 September 15; accepted 2006 November 1; published 2006 November 30

### ABSTRACT

We present spectroscopy of a transit of the exoplanet HD 189733b. By modeling the Rossiter-McLaughlin effect (the anomalous Doppler shift due to the partial eclipse of the rotating stellar surface), we find the angle between the sky projections of the stellar spin axis and orbit normal to be  $\lambda = -1.4 \pm 1.1^\circ$ . This is the third case of a “hot Jupiter” for which  $\lambda$  has been measured. In all three cases  $\lambda$  is small, ruling out random orientations with 99.96% confidence, and suggesting that the inward migration of hot Jupiters generally preserves spin-orbit alignment.

*Subject headings:* planetary systems — planetary systems: formation — stars: individual (HD 189733) — stars: rotation

### 1. INTRODUCTION

A primary reason to study planets of other stars is to learn how typical (or unusual) are the properties of the solar system. For example, the nearly circular orbits of solar system planets were once considered normal, but we now know that eccentric orbits of Jovian planets are common (see, e.g., Halbwachs et al. 2005 or Fig. 3 of Marcy et al. 2005). Likewise, gas giants were once thought to inhabit only the far reaches of planetary systems, an assumption that was exploded by the discovery of “hot Jupiters” (Mayor & Queloz 1995; Butler et al. 1997). This inspired theoretical work on planetary migration mechanisms that can deliver Jovian planets to such tight orbits (as recently reviewed by Thommes & Lissauer [2005] and Papaloizou & Terquem [2006]).

Another striking pattern in the solar system is the close alignment between the planetary orbits and the solar spin axis. The orbit normals of the eight planets are within a few degrees of one another (Cox 2000, p. 295), and the Earth’s orbit normal is only  $7^\circ$  from the solar spin axis (Beck & Giles 2005 and references therein). Presumably this alignment dates back 5 Gyr, when the Sun and planets condensed from a single spinning disk. Whether or not this degree of alignment is universal is unknown. For hot Jupiters in particular, one might wonder whether migration enforces or perturbs spin-orbit alignment.

For exoplanets, the angle between the stellar spin axis and planetary orbit normal (as projected on the sky) can be measured via the Rossiter-McLaughlin (RM) effect: the spectral distortion

observed during a transit due to stellar rotation. The planet hides some of the velocity components that usually contribute to line broadening, resulting in an “anomalous Doppler shift” (Ohta et al. 2005; Giménez 2006; Gaudi & Winn 2007).

Observations of the exoplanetary RM effect have been reported for HD 209458 (Bundy & Marcy 2000; Queloz et al. 2000; Winn et al. 2005; Wittenmyer et al. 2005) and HD 149026 (Wolf et al. 2007). Here we report observations of the RM effect for HD 189733. This system, discovered by Bouchy et al. (2005), consists of a K dwarf with a transiting Jovian planet ( $M_p = 1.15M_J$ ) in a 2.2 day orbit. Our observations are presented in § 2, our model in § 3, and our results in § 4, followed by a discussion in § 5.

### 2. OBSERVATIONS

We observed the transit of UT 2006 August 21 with the Keck I 10 m telescope and the High Resolution Echelle Spectrometer (HIRES; Vogt et al. 1994) following the usual protocols of the California-Carnegie planet search, as summarized here. We employed the red cross-disperser and placed the  $I_2$  absorption cell into the light path to calibrate the instrumental response and the wavelength scale. The slit width was  $0.85''$  and the typical exposure time was 3 minutes, giving a resolution of 70,000 and a signal-to-noise ratio (S/N) of  $300 \text{ pixel}^{-1}$ . We obtained 70 spectra over 7.5 hr bracketing the predicted transit midpoint. To these were added 16 spectra that had been obtained by the California-Carnegie group at random orbital phases.

We determined the relative Doppler shifts with the algorithm of Butler et al. (1996). We estimated the measurement uncertainties based on the scatter in the solutions for each  $2 \text{ \AA}$  section of the spectrum. For the spectra obtained on 2006 August 21 the typical measurement error was  $0.8 \text{ m s}^{-1}$ , while for the other 16 spectra the error was  $\approx 1.3 \text{ m s}^{-1}$ . The data are given in Table 1 and plotted in Figure 1, with enlarged error bars to account for the intrinsic velocity noise of the star (see § 3).

We also needed accurate photometry to pin down the planetary and stellar radii and the orbital inclination. We observed the transit of UT 2006 July 21 with KeplerCam on the 1.2 m telescope at the Fred L. Whipple Observatory on Mt. Hopkins, Arizona. We used the Sloan Digital Sky Survey  $z$ -band filter and an exposure time of 5 s. After bias subtraction and flat-field division, we performed aperture photometry of HD 189733 and 14 comparison stars. The light curve of each com-

<sup>1</sup> Data presented herein were obtained at the W. M. Keck Observatory, which is operated as a scientific partnership among the California Institute of Technology, the University of California, and NASA, and was made possible by the generous financial support of the W. M. Keck Foundation.

<sup>2</sup> Department of Physics, and Kavli Institute for Astrophysics and Space Research, Massachusetts Institute of Technology, Cambridge, MA 02139.

<sup>3</sup> Department of Astronomy, University of California, MC 3411, Berkeley, CA 94720.

<sup>4</sup> Department of Terrestrial Magnetism, Carnegie Institution of Washington, 5241 Broad Branch Road, NW, Washington, DC 20015-1305.

<sup>5</sup> UCO/Lick Observatory, University of California at Santa Cruz, Santa Cruz, CA 95064.

<sup>6</sup> Center of Excellence in Information Systems, Tennessee State University, 3500 John A. Merritt Boulevard, Box 9501, Nashville, TN 37209.

<sup>7</sup> Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138.

<sup>8</sup> Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency, 3-1-1 Yoshinodai, Sagami-hara, Kanagawa 229-8510, Japan.

<sup>9</sup> Department of Physics, University of Tokyo, Tokyo 113-0033, Japan.

<sup>10</sup> Princeton University Observatory, Peyton Hall, Princeton, NJ 08544.

TABLE 1  
RADIAL VELOCITIES OF HD 189733

JD	Radial Velocity (m s <sup>-1</sup> )	Measurement Uncertainty (m s <sup>-1</sup> )	JD	Radial Velocity (m s <sup>-1</sup> )	Measurement Uncertainty (m s <sup>-1</sup> )
2,452,832.881794	-13.964	1.542	2,453,968.807373	45.933	0.816
2,452,898.800937	186.717	1.480	2,453,968.810000	47.080	0.857
2,453,180.918877	153.226	1.609	2,453,968.812650	49.130	0.870
2,453,240.898507	121.542	1.139	2,453,968.815289	45.655	0.982
2,453,303.750382	-202.928	1.244	2,453,968.818021	42.260	0.994
2,453,303.753576	-203.187	1.139	2,453,968.820718	34.659	0.965
2,453,551.963519	-126.301	1.205	2,453,968.823391	27.992	0.987
2,453,693.688738	15.243	1.228	2,453,968.826134	18.132	0.905
2,453,694.690856	-84.160	1.180	2,453,968.828912	7.218	0.858
2,453,695.688148	125.535	1.259	2,453,968.831806	-0.572	1.007
2,453,927.929914	-206.994	1.302	2,453,968.834687	-5.327	0.846
2,453,723.712604	-2.524	1.217	2,453,968.839641	-24.613	0.751
2,453,926.035150	159.489	1.193	2,453,968.842905	-34.630	0.885
2,453,926.887766	-141.263	1.107	2,453,968.845845	-41.573	0.935
2,453,927.929745	53.300	1.157	2,453,968.848877	-45.183	0.842
2,453,934.845208	179.802	1.224	2,453,968.851991	-47.586	0.972
2,453,968.722037	57.606	0.787	2,453,968.855220	-47.746	1.009
2,453,968.724225	57.726	0.802	2,453,968.858356	-44.215	0.928
2,453,968.726447	56.423	0.832	2,453,968.861354	-37.956	0.820
2,453,968.728750	56.045	0.755	2,453,968.864352	-30.035	0.943
2,453,968.731227	54.711	0.848	2,453,968.867361	-27.041	0.890
2,453,968.733611	52.911	0.798	2,453,968.870301	-25.215	0.985
2,453,968.736007	50.628	0.789	2,453,968.873183	-24.080	0.961
2,453,968.757928	42.772	0.726	2,453,968.876076	-22.205	0.892
2,453,968.760266	38.559	0.760	2,453,968.878912	-27.297	0.866
2,453,968.762639	36.305	0.746	2,453,968.881782	-28.557	0.907
2,453,968.765023	37.873	0.741	2,453,968.884583	-29.194	0.822
2,453,968.767419	34.160	0.818	2,453,968.887569	-33.236	0.807
2,453,968.769826	33.982	0.768	2,453,968.890498	-33.070	0.638
2,453,968.772280	32.973	0.657	2,453,968.893403	-31.439	0.847
2,453,968.774757	30.196	0.740	2,453,968.896331	-36.218	0.712
2,453,968.777245	29.968	0.662	2,453,968.899271	-39.051	0.757
2,453,968.779688	28.832	0.651	2,453,968.902118	-41.663	0.900
2,453,968.782095	28.517	0.764	2,453,968.904931	-42.919	0.748
2,453,968.784525	27.562	0.772	2,453,968.907639	-45.150	0.876
2,453,968.786979	26.713	0.682	2,453,968.910278	-42.636	0.956
2,453,968.789433	25.029	0.682	2,453,968.912940	-43.977	0.818
2,453,968.791910	24.544	0.749	2,453,968.915637	-47.332	0.828
2,453,968.794433	20.275	0.661	2,453,968.933565	-58.550	0.715
2,453,968.796944	21.667	0.855	2,453,968.955347	-69.951	0.771
2,453,968.799456	28.013	0.803	2,453,968.977743	-79.481	0.754
2,453,968.802049	31.972	0.780	2,453,969.002674	-95.622	0.750
2,453,968.804745	39.982	0.797	2,453,969.032870	-112.396	0.898

NOTES.— Col. (1) Julian Date at the time of the photon-weighted midexposure. Col. (3) Measurement uncertainties, *not* including photospheric jitter (see § 3).

parison star was normalized to have unit median, and the mean of these normalized light curves was taken to be the comparison signal. The light curve of HD 189733 was divided by the comparison signal and corrected for residual systematic effects by dividing out a linear function based on the out-of-transit data. The light curve is plotted in the top panel of Figure 1.

### 3. THE MODEL

We fitted the fluxes and radial velocities with a parameterized model based on a star and planet in a circular orbit about the center of mass. To calculate the relative flux as a function of the projected separation of the planet and the star, we assumed the limb darkening law to be quadratic and employed the analytic formulas of Mandel & Agol (2002) to compute the integral of the intensity over the unobscured portion of the stellar disk. We fixed the limb-darkening coefficients at the values  $u_1 = 0.320$ ,  $u_2 = 0.267$ , based on the calculations of Claret (2004).

To calculate the anomalous Doppler shift, we used the technique of Winn et al. (2005): we simulated RM spectra with the same data format and noise characteristics as the actual data and determined the Doppler shifts using the same algorithm used on the

actual data. The simulations were based on a “template” spectrum representing the emergent spectrum from a small portion of the photosphere. We scaled the template spectrum in flux by  $\epsilon$  and shifted it in velocity by  $v_p$ , representing the spectrum of the occulted portion of the stellar disk. We subtracted this spectrum from a rotationally broadened version of the template spectrum (broadened to 3 km s<sup>-1</sup> to mimic the disk-integrated spectrum of HD 189733) and then “measured” the anomalous Doppler shift  $\Delta v$ . This was repeated for a grid of  $\{\epsilon, v_p\}$ , and a polynomial function was fitted to the resulting surface.

The template spectrum should be similar to that of HD 189733 but without significant rotational broadening. We tried three different choices: two empirical spectra and one theoretical spectrum. The two empirical spectra were Keck HIRES spectra (S/N  $\approx$  800,  $R \approx 10^5$ ) of HD 3561 (G3 V,  $v \sin i_s = 1.2 \pm 0.5$  km s<sup>-1</sup>) and HD 3765 (K2 V,  $0.0 \pm 0.5$  km s<sup>-1</sup>). The former is 200 K hotter than HD 189733, while the latter is more metal-rich. The theoretical spectrum, with a resolution of 250,000, was taken from Coelho et al. (2005) for a nonrotating star with  $T_{\text{eff}} = 5000$  K,  $\log g = 4.5$ , and  $[\text{Fe}/\text{H}] = 0.0$ .

For each choice of the template spectrum, we derived the




FIG. 1.—Photometry and spectroscopy of HD 189733. *Top*:  $z$ -band photometry during a transit, along with the best-fitting model (*solid line*). *Middle*: Radial velocities as a function of orbital phase (expressed in days), along with the model (*solid line*). *Bottom*: Close-up near the midtransit time. In all cases, the residuals (observed–calculated) are plotted beneath the data.

relation between  $\Delta v$ ,  $\epsilon$ , and  $v_p$ , and optimized the model as described below. With one exception, the results did not depend significantly on the choice of template spectrum (in the sense that measurement errors caused much larger uncertainties). The single exception was  $v \sin i_s$ , for which the results differed as much as 3%. For our final analysis, we used the relation

$$\Delta v = -\epsilon v_p \left[ 1.252 - 0.351 \left( \frac{v_p}{3 \text{ km s}^{-1}} \right)^2 \right] \quad (1)$$

derived from the empirical templates, but we also included an extra error term of 6% in  $v \sin i_s$  as a conservative estimate of the systematic error. In summary, the projected separation of the planet and the star determines the transit depth  $\epsilon$  and the subplanet rotation velocity<sup>11</sup>  $v_p$ , and then equation (1) is used to calculate the anomalous Doppler shift.

<sup>11</sup> The subplanet velocity is the projected rotation velocity of the portion of the star hidden by the planet, and is calculated assuming no differential rotation, an assumption justified by Gaudi & Winn (2007).

The fitting statistic was

$$\chi^2 = \sum_{j=1}^{86} \left[ \frac{v_j(\text{obs}) - v_j(\text{calc})}{\sigma_{v,j}} \right]^2 + \sum_{j=1}^{752} \left[ \frac{f_j(\text{obs}) - f_j(\text{calc})}{\sigma_{f,j}} \right]^2 + \left( \frac{\Delta\gamma}{12 \text{ m s}^{-1}} \right)^2 + \left( \frac{M_s/M_\odot - 0.82}{0.03} \right)^2, \quad (2)$$

where  $f_j(\text{obs})$  is the flux observed at time  $j$ ,  $\sigma_{f,j}$  is the corresponding uncertainty, and  $f_j(\text{calc})$  is the calculated value. A similar notation applies to the velocities. The last two terms are a priori constraints explained below.

It is important for  $\sigma_{f,j}$  and  $\sigma_{v,j}$  to include not only measurement errors but also any unmodeled systematic errors. To account for systematic errors in the photometry, we increased the Poisson estimates of the errors by a factor of 1.2, at which point  $\chi^2/N_{\text{DOF}} = 1$  when fitting only the fluxes. Determining the appropriate weights for the velocities was more complex. HD 189733 is chromospherically active and should exhibit velocity noise (“photospheric jitter”) with an amplitude of  $11 \text{ m s}^{-1}$  according to the empirical relations of Wright (2005). However, the timescale of the noise cannot be predicted as easily. Noise from spots or plages would occur on the timescale of the rotation period ( $\approx 13$  days), while noise from oscillations and flows occurs on shorter timescales.

We took the following approach. First, we fitted only the 16 velocities obtained sporadically prior to 2006 August 21 and found the rms residual to be  $12 \text{ m s}^{-1}$ , in agreement with the Wright (2005) relations. Therefore, for fitting purposes, we inflated the error bars  $\sigma_{v,j}$  of those 16 velocities to  $12 \text{ m s}^{-1}$ . Second, we fitted only the 44 “out-of-transit” velocities measured on 2006 August 21 and found the rms residual to be  $1.5 \text{ m s}^{-1}$ . In addition, there were correlations in the residuals on a timescale of  $\sim 15$  minutes ( $\sim 4$  data points). The correlations effectively reduce the number of independent data points by 4, or equivalently, they double the error per point. Therefore, for fitting purposes, we inflated the error bars  $\sigma_{v,j}$  of all the 2006 August 21 velocities to  $3 \text{ m s}^{-1}$ . Apparently, for HD 189733, most of the velocity noise occurs on a timescale longer than one night.

Our free parameters were the two bodies’ masses and radii ( $M_s$ ,  $M_p$ ,  $R_s$ , and  $R_p$ ), the orbital inclination ( $i$ ), the midtransit time ( $T_c$ ), the line-of-sight stellar rotation velocity ( $v \sin i_s$ ), the angle between the projected stellar spin axis and orbit normal ( $\lambda$ ; see Ohta et al. 2005 or Gaudi & Winn 2007 for a diagram of the coordinate system), the velocity zero point ( $\gamma$ ), a velocity offset specific to the night of 2006 August 21 ( $\Delta\gamma$ ), and a long-term velocity gradient  $\dot{\gamma}$ . The parameter  $\Delta\gamma$  is needed because of the photospheric jitter; the first a priori constraint in equation (2) enforces a reasonable level of noise. The gradient  $\dot{\gamma}$  was included to account for the long-period orbit of HD 189733 and its companion star (Bakos et al. 2006a) or possible long-period planets. We fixed the orbital period to be 2.218575 days (Bouchy et al. 2005; Hébrard & Lecavelier Des Etangs 2006). A well-known degeneracy prevents  $M_s$ ,  $R_s$ , and  $R_p$  from being determined independently. We broke this degeneracy with the second a priori constraint in equation (2), which enforces the spectroscopic determination of  $M_s$  by Bouchy et al. (2005).

We used a Markov Chain Monte Carlo algorithm to solve for the model parameters and their uncertainties (see, e.g., Tegmark et al. 2004 or Ford 2005). Our jump function was the addition of a Gaussian random number to each parameter value. We set the perturbation sizes such that  $\sim 20\%$  of jumps are executed. We created 10 independent chains, each with 500,000

TABLE 2  
SYSTEM PARAMETERS OF HD 189733

Parameter	Value	Uncertainty
$M_S$ ( $M_\odot$ )	0.82	0.03
$M_p$ ( $M_J$ )	1.13	0.03
$R_S$ ( $R_\odot$ )	0.73	0.02
$R_p$ ( $R_J$ )	1.10	0.03
$i$ (deg)	86.1	0.2
$T_c$ (HJD)	2,453,937.7759	0.0001
$v \sin i_s$ ( $\text{km s}^{-1}$ )	2.97	0.22
$\lambda$ (deg)	-1.4	1.1
$\gamma$ ( $\text{m s}^{-1}$ )	5.0	10.1
$\Delta\gamma$ ( $\text{m s}^{-1}$ )	-15.0	4.8
$\dot{\gamma}$ ( $\text{m s}^{-1} \text{yr}^{-1}$ )	-1.9	3.3

points, starting from random initial positions, and discarded the first 20% of the points in each chain. The Gelman & Rubin (1992)  $R$  statistic was close to unity for each parameter, a sign of good mixing and convergence. We merged the chains and took the median value of each parameter to be our best estimate, and the standard deviation as the  $1 \sigma$  uncertainty. For the special case of  $v \sin i_s$ , we added an additional error of 6% in quadrature, due to the systematic error noted previously.

#### 4. RESULTS

The results are given in Table 2. Those parameters depending chiefly on the photometry ( $R_p$ ,  $R_s$ ,  $i$ ) are in agreement with the most accurate results reported previously (Bakos et al. 2006b). Likewise, our result for the planetary mass,  $M_p = 1.13M_J \pm 0.03M_J$ , agrees with the value  $1.15M_J \pm 0.04M_J$  measured by Bouchy et al. (2005). Our result for the projected rotation velocity is  $v \sin i_s = 3.0 \pm 0.2 \text{ km s}^{-1}$ . This agrees with the value  $3.5 \pm 1.0 \text{ km s}^{-1}$  reported by Bouchy et al. (2005), which was based on the line broadening in their disk-integrated stellar spectrum. It also agrees with the value  $3.2 \pm 0.7 \text{ km s}^{-1}$  based on a similar analysis of our own Keck spectra (D. Fischer 2006, private communication). The most interesting result is  $\lambda = -1.4 \pm 1.1$ . The sky projections of the stellar spin axis and the orbit normal are aligned to within a few degrees.

#### 5. DISCUSSION

HD 189733 is the third exoplanetary system (and the third hot Jupiter) for which it has been possible to measure  $\lambda$ . The

first system was HD 209458, for which  $\lambda = -4.4^\circ \pm 1.4^\circ$  (Winn et al. 2005; see also Wittenmyer et al. 2005, who modeled the RM effect but required  $\lambda = 0$ ). The second system was HD 149026 (Wolf et al. 2007), for which  $\lambda = 11^\circ \pm 14^\circ$ . The small observed values of  $\lambda$  suggest that the most common end-state of the inward migration of a hot Jupiter involves a close alignment.

With only three systems, we cannot yet measure the distribution of  $\lambda$ , but we can test the hypothesis of random orientations (i.e., a uniform distribution in  $\lambda$ ). The weighted mean of the measured values of  $|\lambda|$  is  $2.6^\circ$ . If we replace the measured values by random numbers drawn from a uniform distribution, the probability that the weighted mean<sup>12</sup> will be this small is only 0.04%. Hence, we rule out the hypothesis of random orientations with 99.96% confidence.

Winn et al. (2005) argued that tides from the star would not ordinarily cause alignment within the star's main-sequence lifetime. There are thus two basic possibilities: either the alignment is primordial and was not disturbed by migration, or there was a different mechanism to damp any initial or induced misalignment. Among the various theories of hot Jupiter migration, some would tend to enhance any initial misalignments and are thereby constrained by our results. Such scenarios include planet-planet scattering followed by circularization (Rasio & Ford 1996; Weidenschilling & Marzari 1996), Kozai migration (Wu & Murray 2003; Eggenberger et al. 2004), and tidal capture (Gaudi 2003).

The agreement among the three systems studied to date is clear but should not discourage future measurements. Obviously, a sample of three is only barely sufficient to draw a conclusion. And of course, the discovery of even a single example of a grossly misaligned system would be of great interest.

We thank Debra Fischer for running SME on our spectra and Scott Gaudi for very helpful discussions. We recognize and acknowledge the very significant cultural role and reverence that the summit of Mauna Kea has always had within the indigenous Hawaiian community. We are most fortunate to have the opportunity to conduct observations from this mountain.

<sup>12</sup> Here we have assumed the uncertainty in  $\lambda$  is independent of  $\lambda$ , a good approximation because all three systems have an intermediate impact parameter (Gaudi & Winn 2007).

#### REFERENCES

- Bakos, G. Á., Pál, A., Latham, D. W., Noyes, R. W., & Stefanik, R. P. 2006a, *ApJ*, 641, L57  
 Bakos, G. Á., et al. 2006b, *ApJ*, submitted (astro-ph/0603291)  
 Beck, J. G., & Giles, P. 2005, *ApJ*, 621, L153  
 Bouchy, F., et al. 2005, *A&A*, 444, L15  
 Bundy, K. A., & Marcy, G. W. 2000, *PASP*, 112, 1421  
 Butler, R. P., Marcy, G. W., Williams, E., McCarthy, C., Dosanjh, P., & Vogt, S. S. 1996, *PASP*, 108, 500  
 Butler, R. P., Marcy, G. W., Williams, E., Hauser, H., & Shirts, P. 1997, *ApJ*, 474, L115  
 Claret, A. 2004, *A&A*, 428, 1001  
 Coelho, P., Barbuy, B., Meléndez, J., Schiavon, R. P., & Castilho, B. V. 2005, *A&A*, 443, 735  
 Cox, A. N. 2000, *Allen's Astrophysical Quantities* (4th ed.; New York: AIP)  
 Eggenberger, A., Udry, S., & Mayor, M. 2004, *A&A*, 417, 353  
 Ford, E. B. 2005, *AJ*, 129, 1706  
 Gaudi, B. S. 2003, *ApJ*, submitted (astro-ph/0307280)  
 Gaudi, B. S., & Winn, J. N. 2007, *ApJ*, in press (astro-ph/0608071)  
 Gelman, A., & Rubin, D. B. 1992, *Stat. Sci.*, 7, 457  
 Giménez, A. 2006, *ApJ*, 650, 408  
 Halbwachs, J. L., Mayor, M., & Udry, S. 2005, *A&A*, 431, 1129  
 Hébrard, G., & Lecavelier Des Etangs, A. 2006, *A&A*, 445, 341  
 Mandel, K., & Agol, E. 2002, *ApJ*, 580, L171  
 Marcy, G., Butler, R. P., Fischer, D., Vogt, S., Wright, J. T., Tinney, C. G., & Jones, H. R. A. 2005, *Prog. Theor. Phys. Suppl.*, 158, 24  
 Mayor, M., & Queloz, D. 1995, *Nature*, 378, 355  
 Ohta, Y., Taruya, A., & Suto, Y. 2005, *ApJ*, 622, 1118  
 Papaloizou, J. C. B., & Terquem, C. 2006, *Rep. Prog. Phys.*, 69, 119  
 Queloz, D., Eggenberger, A., Mayor, M., Perrier, C., Beuzit, J. L., Naef, D., Sivan, J. P., & Udry, S. 2000, *A&A*, 359, L13  
 Rasio, F. A., & Ford, E. B. 1996, *Science*, 274, 954  
 Tegmark, M., et al. 2004, *Phys. Rev. D*, 69, 103501  
 Thommes, E. W., & Lissauer, J. J. 2005, in *Astrophysics of Life*, ed. M. Livio, N. Reid, & W. Sparks (New York: Cambridge Univ. Press), 41  
 Vogt, S. S., et al. 1994, *Proc. SPIE*, 2198, 362  
 Weidenschilling, S. J., & Marzari, F. 1996, *Nature*, 384, 619  
 Winn, J. N., et al. 2005, *ApJ*, 631, 1215  
 Wittenmyer, R. A., et al. 2005, *ApJ*, 632, 1157  
 Wolf, A., et al. 2007, *ApJ*, in press  
 Wright, J. T. 2005, *PASP*, 117, 657  
 Wu, Y., & Murray, N. 2003, *ApJ*, 589, 605